


In our experiment⁴ four ZnSe nanoparticles samples, one with low Mn²⁺ concentration (A), one with an intermediate Mn²⁺ concentration (B), one with a high Mn²⁺ concentration (C), and one with no Mn²⁺, were synthesized. The sample with no Mn²⁺ had a sharp ZnSe bandedge emission peak and a quantum yield of ~29%. The samples with Mn²⁺ had a significant decrease in bandedge emission. Sample A had no Mn²⁺ $^4T_1 \rightarrow ^6A_1$ emission, but showed some ZnSe bandedge emission and trap state emission. Sample B had Mn²⁺ $^4T_1 \rightarrow ^6A_1$ emission and a further reduction in ZnSe bandedge emission and trap state emission. Sample C showed an increase in the Mn²⁺ $^4T_1 \rightarrow ^6A_1$ emission, a dramatic increase in trap state emission, and essentially no ZnSe bandedge emission. In order to better understand these observations XAFS data were taken. The XAFS indicated that there was a reduction in the Zn and Mn first neighbor Se coordination from the bulk value but a lack of a reduction in the Se first neighbor coordination. This suggests that the core of the nanoparticles resembles that of bulk ZnSe, and the surface of the particle has a higher concentration of metal atoms. We propose that the surface Mn²⁺ possessed an octahedral geometry, and the overall low emission quantum yield is primarily due to the presence of Mn²⁺ on the particle surface.

- (1) Hines, M. A.; Guyot-Sionnest, P. *J. Phys. Chem. B* **1998**, *102*, 3655.
- (2) Suyver, J. F.; Wuister, S. F.; Kelly, J. J.; Meijerink, A. *Phys. Chem. Chem. Phys.* **2000**, *2*, 5445.
- (3) Norris, D. J.; Nan, Y.; Charnock, F. T.; Kennedy, T. A. *Nano Lett.* **2001**, *1*, 3.
- (4) Norman Jr., T. J.; Magana, D.; Wilson, T.; Burns, C.; Zhang, J. Z.; Cao, D.; Bridges, F. *J. Phys. Chem. B* **2003**, *107*, 6309.



References

(9) Norris, D. J.; Nan, Y.; Charnock, F. T.; Kennedy, T. A. *Nano Lett.* 2001, 1, 3-7

(10) Borse, P. H.; Srinivas, D.; Shinde, R. F.; Date, S. K.; Vogel, W.; Kulkarni, S. K. *Phys. Rev. B* 1999, 60, 8659-8664

(11) Courio, G.; Esnouf, S.; Gacoin, T.; Boilot, J. P. *J. Chem. Phys.* 1996, 100, 20021.

